Proceedings of the 19th

SUNFLOWER RESEARCH WORKSHOP

January 9 & 10, 1997

Sponsored by:
National
Sunflower
Association

The 1996 Sunflower Research Workshop, sponsored by the National Sunflower Association, took place on January 9 and 10, 1997, at the Ramada Plaza Suites, Fargo, ND. The workshop was very well attended and received by public and private researchers from the United States and Canada, as well as other interested parties.

This volume contains nearly all the presentations given at the 1996 workshop. Some of the papers are summarized or abstract form.

The National Sunflower Association would like to extend its appreciation to those presenting papers/posters at this annual Sunflower Research Workshop and to those who participated by their

attendance and questions. Special thanks are extended to the NSA Research Forum Planning Committee, Dr. Mary Lou Straley, Cargill, Inc.; Dr. Brady Vick and Dr. Laurence D. Charlet, USDA-ARS. Thanks also to Steve Sebesta, Cargill Hybrid Seeds; Gary Brewer, NDSU Entomology; and Brady Vick, USDA-ARS Oilseeds Research for their expertise in moderating the workshop sessions.

Questions regarding these proceedings may be directed to the National Sunflower Association, 4023 State Street, Bismarck, ND 58501.

Note: The papers in these proceedings should not be reprinted in part or in total without the expressed consent of the author(s) involved.

TABLE OF CONTENTS

BREEDING AND GENETICS	ENTOMOLOGY
Cytoplasmic Male Sterility and Fertility	Overwintering Biology of the Sunflower Beetle,
Restoration in a Helianthus Annuus	Zygogramma Exclamationis (Coleoptera:
Landrace PI4325131	Chrysomelidae), and the Impact of Tillage on
C. C. Jan and Brady A. Vick	Survival and Emergence29
	Larry Charlet
Influence of Parental Selection on Mid-Oleic	
Acid Levels of Sunflower Oil3	Control of Sunflower Stem Weevil with Planting
J. F. Miller and B. A. Vick	Time Treatments in Dryland Sunflowers,
	199635
Wild Sunflowers as a Potential Source of	Scott Armstrong and Mike Koch
Disease Resistance for Cultivated Sunflower8	
Gerald J. Seiler	Overview of Factors Involved in Banded
	Sunflower Moth Oviposition on Sunflower37
Development of High Linoleic Sunflower	John F. Barker
Hybrids for the U.S14	3
Jerry F. Miller and Brady A. Vick	Red Sunflower Seed Weevil Outlook and
	Management41
	Gary L. Brewer and Phil Glogoza
BIOTECH AND OILSEEDS TRADE	,
World Trade in Minor Oilseeds: Impacts of the	Sunflower Midge - A Proposal for Future
Changing Policy Environment17	Research43
Demcey Johnson and Frank Dooley	Gary J. Brewer and Larry Charlet
Molecular Markers Associated with Vigor	Acquisition and Application of Bacillus
Restoration from Perennial Sunflower	thuringlensis to Sunflower by Honeybees49
Species26	J. L. Jyoti and Gary J. Brewer
W. Scott Grayburn, C. C. Jan and	<i>y</i> = <i>y</i> , = = = = = = , <i>y</i> = = = = =
Brady A. Vick	PRODUCTION
Diady II. VICE	Micro Nutrient Fertilizer52
Regulatory Requirements for the Field Testing	Tom Johnson
of Genetically Engineered Sunflowers28	2 om Johnson
Arnold Foudin	Cultural Systems for Reduced Pesticide Use in
Amoid Foudin	Sunflower55
	Donald L. Tanaka and Randy L. Anderson
	Donald L. Tallaka and Randy L. Anderson

Narrow Row Seeding of Non-Oll Sunflowers	Progress on Reducing Blackbird Damage to Sunflower: Status of Avicide Research
PATHOLOGY Inoculation Methods for the Assessment of Sclerotinia Headrot in Sunflower72 Khalid Rashid	M. L. Avery, and W. J. Bleier Presentation of the Advanta Sunflower RFLP Linkage Map for Public Research
Phomopsis Stem Canker Resistance in USDA and Commerical Sunflower Germplasm76 T. J. Gulya	Comparative Assay of Sunflower Beetle Preferences for Twelve Commerical Hybrids119 Craig R. Roseland and Teryl J. Grosz
Central Great Plains Sunflower Rust Situation in 1996: Race Identification and Hybrid Evaluation for Resistance	
Sunflower Diseases in South Africa in 1996	
Screening for Resistance in Sunflower Against Albugo Tragopogonls	
Isozyme Genetic Markers and Helianthus Germplasm Management89 R. C. Cronn, M.E. Brothers, K. M. Klier, P. K. Bretting, and J. F. Wendel	
Interspecific Hybrids Between Cultivated Sunflower Susceptible to Orobanche Cernua and the Wild Resistant Species H. Laevigatus and H. Giganteus97 S. Sukno, C. C. Jan, J. Melero-Vara, and J. M. Fernandez-Martinez	
Predicting Sunflower Stem Weevil Emergence on the Central Great Plains Using Degree Days107	

J. Scott Armstrong and Mike D. Koch

PROGRESS ON IDENTIFYING THE NONTARGET HAZARDS ASSOCIATED WITH THE USE OF DRC-1339 A VICIDE IN SOUTH DAKOTA

G. A. KNUTSEN¹, M. J. KENYON¹, G. M. LINZ², M. L. AVERY³, AND W. J. BLEIER¹

¹Department of Zoology, North Dakota State University, Fargo, ND 58105, U.S.A.

²USDA-APHIS Great Plains Field Station, Bismarck, ND 58504, U.S.A.

³USDA-APHIS Florida Field Station, Gainesville, FL 32641, U.S.A.

In August and September, 25 million blackbirds migrate through sunflower growing areas of North Dakota, South Dakota, and Minnesota. Sunflower growers forfeit about \$5.0 million to these birds. Moreover, the cost of resources (e.g., vehicle, ammunition, scare devices and people) needed to protect fields from blackbirds is expensive. Growers and government agencies are using non-lethal techniques to reduce blackbird damage. However, methods developed thus far have limitations of costs, logistics and effectiveness.

In 1995, after six years of research in Louisiana and Texas, the Environmental Protection Agency (EPA) granted a Section 3 label for 'Compound DRC-1339 (3-chloro-4-methylbenzenamine HCL) Concentrate-Staging Areas' for bird damage control. A single seed of 2% (wt/wt) treated rice is highly toxic to blackbirds but many nontarget birds either are resistant to the chemical (e.g., house sparrows, birds-of-prey) or must ingest many treated seeds (e.g., ring-necked pheasants, waterfowl) to acquire a lethal dose of the avicide. Because acquiring a lethal dose of DRC-1339 is possible for pheasant and waterfowl, officials of state and federal agencies have emphasized that suitable testing must be completed before the operational use of DRC-1339 is started in the northern Great Plains.

Since 1993, quantitative data on the nontarget hazards of using DRC-1339 -treated rice in harvested grain fields in east-central South Dakota have been collected. In March and April 1993-1996, harvested corn and soybean fields were baited with untreated brown rice and brown rice treated with DRC-1339 avicide. Nontarget birds appeared to use corn stubble fields more than soybean fields. Of the 32 bird species visiting these fields, mallards, Canada geese, western meadowlarks, American tree sparrows, and killdeer were the most commonly observed species. The ring-necked pheasant and western meadowlark may be at the highest risk for acquiring a lethal dose of DRC-1339 because of their feeding ecology and susceptability to DRC-1339.

We evaluated the risk of avicide baiting to ring-necked pheasants at the Florida Field Station of the National Wildlife Research Center. In one study, 10 of 12 individually caged female pheasants preferred to eat cracked corn and sorghum over rice. In another study, brown rice eaten by pheasants over four days, while penned in 0.2 ha enclosure, exceeded that needed to acquire a median acute lethal dose of DRC-1339.

Based on consultations with various federal and state agencies, questions related to the primary hazards of DRC-1339 to nontarget birds, particularly pheasants, must be fully assessed. In 1997, retriever dogs will be tested for augmenting searches for dead birds, the number and kinds of wildlife scavenging on dead blackbirds will be documented, and specialized video equipment will be used for assessing nontarget hazards in harvested fields. A radio-telemetry study documenting the effects of DRC-1339-treated rice baits on the abundance of pheasants in the baited area will be conducted by South Dakota State University in 1998 and 1999.